



1/26




FIG. 1A

2/26





FIG. 1B


3/26

FIGURE 2A




4/26

FIGURE 2B




5/26

FIG. 3A




6/26

FIG. 3B




7/26

FIG. 4



8/26


FIGURE 5






9/26

FIGURE 6A




10/26


FIGURE 6B



11/26


FIGURE 7






13/26

**FIG. 8C**



14/26

FIG. 9A



**FIG. 9B**





Figure 1 is a line graph showing the reciprocal titers of IgM ELISA for various antigens over a 36-week period. The Y-axis represents 'IgM ELISA TITERS (RECIPROCAL)' with a scale break between 200 and 5120. The X-axis represents 'WEEKS' from 0 to 36. The graph includes data for three antigens: Cy, GM2-KLH + 200  $\mu$ g QS-21, and 200  $\mu$ g QS-21. Titers generally increase over time, with some peaks around week 16 and week 24.

Week	Cy	GM2-KLH + 200 $\mu$ g QS-21	200 $\mu$ g QS-21
0	0	0	0
4	100	100	100
8	100	100	100
12	100	100	100
16	100	100	100
20	100	100	100
24	100	100	100
28	100	100	100
32	100	100	100
36	100	100	100




17/26

FIG. 9D




18/26

FIG. 10A




19/26

FIG. 10B




20/26


FIGURE 11A



21/26


FIGURE 11B






23/26

FIGURE 13



24/26


**FIGURE 14**





25/26

FIGURE 15



26/26

**FIGURE 16**

